

Gamma Rays of Tellurium-131 and Tellurium-129

C. A. MALLMANN, A. H. W. ATEN, JR.,* D. R. BES, AND CLARA M. DE McMILLAN
Comisión Nacional de la Energía Atómica, Buenos Aires, Argentina

(Received February 28, 1955)

The γ rays of Te^{131} and Te^{129} have been investigated by means of a crystal spectrometer.

EXPERIMENTAL

THE γ rays of Te^{131} and Te^{129} were studied in the laboratory for nuclear spectroscopy of this Institute by means of a crystal spectrometer equipped with a sliding channel. Energy and efficiency calibrations were performed by means of Hg^{203} , Cs^{137} , Na^{22} , and Y^{88} , and only photopeaks were used.

Most of the samples were prepared by irradiation of sodium tellurite with 26-Mev deuterons in the synchrocyclotron of this Institute. After iodine had been extracted twice, tellurium was precipitated in the presence of suitable hold-back carriers, and purified by a scavenging precipitation of ferric hydroxide and a second tellurium precipitation. In some cases Te^{129} was obtained free from Te^{131} via an antimony fraction separated by hydride formation from uranium oxide irradiated with deuterons. (These separations were performed in the Radiochemistry Laboratories.)

TELLURIUM-131

In the 25-minute period of Te^{131} the following γ energies were found: 145 ± 10 kev (100), 450 ± 10 kev (24), 595 ± 15 kev (about 6), 950 ± 15 kev (about 4) and 1140 ± 20 kev (about 8). The figures given in parentheses indicate relative intensities, the last three are only rough approximations. Measurements of the total γ energy by means of a Geiger counter with known γ sensitivity indicate that about one γ ray of 145 kev occurs per β decay, and therefore the intensity ratios also give a rough value of the numbers of each γ ray per β process.

In the 30-hour period of Te^{131} the energies observed were: 145 ± 10 kev, 770 ± 20 kev, and 1140 ± 20 kev. The intensity of the 770 kev is at least about as high as that of the 145 kev, and the 1140 kev is much stronger here relative to the 145-kev line than it is in the 25-minute isomer. Because of the complex nature of the spectrum it was not possible to observe the 450-kev and the 595-kev γ rays here with certainty.

It should be mentioned that our results seem to be in agreement with the experiments of Geiger¹ if it is

* On leave of absence from the Instituut voor Kernfysisch Onderzoek, Amsterdam. Sincere thanks are due the Director of the Comisión Nacional de la Energía Atómica for a generous invitation which made possible this collaboration.

¹ K. Geiger, Z. Naturforsch. **7a**, 579 (1952).

realized that the γ rays of higher energy cannot be recognized separately in absorption measurements.

It seems to be evident that the 145-kev transition goes to the ground state. It is tempting to assume that the 440-kev transition goes to the 145-kev level and that the 595-kev line represents the crossover of these two lines. From the difference in the intensity ratios in the 25-minute period and the 30-hour period, we may conclude that the 1140-kev γ ray and the 950-kev γ ray do not come from the same level. The main fraction of the 25-minute isomer seems to go to the 145-kev level.

In the decay of the 30-hour isomer the principal feature is the high intensity of the 770-kev γ ray, a line which we did not observe in the 25-minute period. Its occurrence is explained most easily by assuming a direct β decay of the metastable state of Te^{131} with a frequency comparable to that of the isomeric transition. Goldhaber and Hill² have given arguments for the possibility of such a direct β transition on the basis of the yield of the chemical separation of the two isomers. A very important fraction if not the main part of the direct β decay of the 30-hour isomer is followed by the 770 kev γ ray, a smaller fraction by the 1140-kev γ ray.

TELLURIUM-129

By far the larger part of the β decay of the 72-minute period of Te^{129} goes to the ground state, unless there should be a low-lying level in I^{129} with an energy below about 70 kev. A γ line of 435 ± 20 kev occurs in about 9 percent of the β transitions, and a γ line of 1080 ± 40 kev in about 0.7 percent.

ACKNOWLEDGMENTS

A fuller discussion of this work will be published in the *Publicaciones de la Comisión Nacional de la Energía Atómica, Argentina*. We wish to thank Ing. E. Galloni and Mr. Monneret de Villars for the cyclotron irradiations, Dr. K. Fraenz and Ing. Pinasco for help with the construction of the crystal spectrometer, and S. Mayo, J. Peyre, R. Norscini, and Dr. H. Bosch for assistance with the measurements.

² M. Goldhaber and R. D. Hill, Revs. Modern Phys. **24**, 179 (1952).